# Stop an Overdose SAVE A LIFE

If you *or* someone you know uses opioids (such as heroin or prescription pain medications), please ask our emergency department staff for information on how to get a lifesaving naloxone kit.

When a person overdoses on opioids, they stop breathing. Naloxone reverses the overdose, allowing the victim to breathe again. It is a safe antidote for opioid overdose, and does not harm the person if they have not taken opioids.

## A person may be having an opioid overdose if they:

- Are known to use opioids
- Will not wake up, even when you shake them
- Are breathing slowly or not at all
- Have lips or fingernails that are blue or gray-colored
- Have clammy or pale skin

### IF YOU SEE THESE SIGNS, IT MAY BE AN OPIOID OVERDOSE!

# **Opioids include:**

GENERIC	BRAND NAME	GENERIC	BRAND NAME
Buprenorphine	Suboxone, Subutex, Zubsolv, Bunavail, Butrans	Hydromorphone	Dilaudid
		Methadone	Dolophine, Methadose
Codeine	Tylenol with codeine, Tylenol #3	Morphine	MSContin, Embeda, Kadian, Avinza
Fentanyl	Duragesic	Oxycodone	Percocet, OxyContin, Percodan, Roxicodone
Hydrocodone	Vicodin, Norco, Lorcet, Lortab,		
,	Zohydro	Oxymorphone	Opana
HEROIN is also an opioid.			

# NALOXONE KITS AVAILABLE HERE.



